

Universal Analytic Gröbner Bases and Tropical Geometry

Tristan Vaccon

Université de Limoges; CNRS, XLIM UMR 7252
Limoges, France
tristan.vaccon@unilim.fr

Thibaut Verron

Johannes Kepler University, Institute for Algebra
Linz, Austria
thibaut.verron@jku.at

ABSTRACT

A universal analytic Gröbner basis (UAGB) of an ideal of a Tate algebra is a set containing a local Gröbner basis for all suitable convergence radii. In a previous article, the authors proved the existence of finite UAGB's for polynomial ideals, leaving open the question of how to compute them. In this paper, we provide an algorithm computing a UAGB for a given polynomial ideal, by traversing the Gröbner fan of the ideal. As an application, it offers a new point of view on algorithms for computing tropical varieties of homogeneous polynomial ideals, which typically rely on lifting the computations to an algebra of power series.

Motivated by effective computations in tropical analytic geometry, we also examine local bases for more general convergence conditions, constraining the radii to a convex polyhedron. In this setting, we provide an algorithm to compute local Gröbner bases and discuss obstacles towards proving the existence of finite UAGBs.

CCS CONCEPTS

• Computing methodologies → Algebraic algorithms.

KEYWORDS

Algorithms, Gröbner bases, Tate algebra, Tropical Geometry, Universal Gröbner basis

ACM Reference Format:

Tristan Vaccon and Thibaut Verron. 2023. Universal Analytic Gröbner Bases and Tropical Geometry. In *International Symposium on Symbolic and Algebraic Computation 2023 (ISSAC 2023)*, July 24–27, 2023, Tromsø, Norway. ACM, New York, NY, USA, 9 pages. <https://doi.org/10.1145/3597066.3597110>

1 INTRODUCTION

The notion of Tate algebras has been introduced by Tate in [25] to develop analytic geometry over the p -adics, founding what is now called rigid geometry. This theory has proved to be central to many developments in number theory. In this context, Tate algebras and ideals in Tate algebras serve the same purpose as polynomial algebras and polynomial ideals in classical algebraic geometry. Tate algebras are defined as algebras of power series over a complete discrete valuation field with convergence conditions such as converging on a given ball or a polydisk with given radii.

T. Verron was supported by the Austrian FWF grant P34872.

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).
ISSAC 2023, July 24–27, 2023, Tromsø, Norway
© 2023 Copyright held by the owner/author(s).
ACM ISBN 979-8-4007-0039-2/23/07.
<https://doi.org/10.1145/3597066.3597110>

In previous works [4–6], the authors showed that it is possible to define and compute Gröbner bases (GB) of ideals in Tate algebras, and modern algorithms for Gröbner bases computations like signature-based algorithms [5] and FGLM [6] can be adapted to this setting.

In [6, 7], the authors paved the way for computations in Tate algebras in case of overconvergence, e.g. ideals defined by series converging on a bigger polydisk. Motivated by the application of analytic geometry in algebraic geometry, an extreme example of this phenomenon is that of ideals defined by polynomials in a Tate algebra. In [7], it was proved that it is possible to compute a GB of a polynomial ideal in a Tate algebra that is made of polynomials. It was also proved that for any polynomial ideal, there exists a universal analytic Gröbner basis (UAGB), i.e. a finite list of polynomials such that whenever they are seen as converging power series in a Tate algebra, they form a Gröbner basis of the corresponding ideal in this algebra. Such a UAGB of a polynomial ideal then contains abundant information on the local behavior of the ideal. In this paper, we prove that a UAGB is also universal irrespectively of the order used as tie-break in the algebra. Furthermore, we provide an algorithm to compute a UAGB in finite time (Algorithm 2, Theorem 4.7).

From a universal GB, it is natural to consider computing the tropical variety of an ideal. Over a field with valuation K such as \mathbb{Q}_p or $\mathbb{F}_p((t))$, the tropical variety $\text{trop}(V)$ of a variety V defined by an ideal I can be defined as the closure of the image of V by the valuation, or alternatively using conditions on leading terms of the elements of I . Acting as a combinatorial shadow of V , many information on V can be recovered from $\text{trop}(V)$. The developments of tropical geometry have been plentiful. To only name a few: enumerative geometry [16], understanding optimization algorithms [1] or analyzing artificial neural networks of the ReLU type.

Universal GB can help in the computation of $\text{trop}(V)$ using the second definition of the tropical variety. In our context, working with Tate algebras instead of polynomial rings gives rise to *tropical analytic geometry*. This emerging field has been defined in [21], adapting the language of tropical geometry to the world of rigid geometry.

In Section 5, we consider the case of tropical varieties of Tate polynomial ideals. We show that the tropical variety of a polynomial ideal is the union of the tropical varieties of its Tate completions, which allows to compute the tropical variety using universal analytical Gröbner bases and the Gröbner fan. This matches what was known for ideals in $k[[T]][X]$, which were used as a lifting target in existing algorithms for computing tropical varieties over valued fields. As such, we provide a new point of view on those algorithms, allowing them to work directly on the Tate series without lifting. To the best of our knowledge, this is the first effective application of tropical analytic geometry.

Finally, motivated by going further into the development of effective computations in rigid geometry, we aim at building up the tools for effective computations on *affinoid subdomains*. Roughly speaking, affinoid subdomains are constructed using generalizations of Tate algebras to more general convergence conditions (e.g. converging on an annulus) and taking quotients by ideals.

In Section 6, we make some steps into this journey by providing effective computations of local GB in the special case of some polyhedral subdomains as defined in [21]. We conclude with some conjectures on UAGB in this context, along with examples and comments.

2 SETTING

2.1 Tate algebras and Gröbner bases

In this section, we recall the definition of Tate algebras and their theory of Gröbner bases (GB for short). Let K be a field with a valuation val making it complete. Let π be a uniformizer of K , that is an element of valuation 1. Typical examples of such a setting are p -adic fields like $K = \mathbb{Q}_p$ with $\pi = p$ or Laurent series fields like $K = \mathbb{Q}((T))$ with $\pi = T$.

For $\mathbf{r} = (r_1, \dots, r_n) \in \mathbb{Q}^n$, the *Tate algebra* $K\{X; \mathbf{r}\}$ is defined as

$$K\{X; \mathbf{r}\} := \left\{ \sum_{\mathbf{i} \in \mathbb{N}^n} a_{\mathbf{i}} X^{\mathbf{i}} \text{ s.t. } a_{\mathbf{i}} \in K \text{ and } \text{val}(a_{\mathbf{i}}) - \mathbf{r} \cdot \mathbf{i} \xrightarrow{|\mathbf{i}| \rightarrow +\infty} +\infty \right\}$$

We call the tuple \mathbf{r} the convergence log-radii of the Tate algebra. We define the Gauss valuation of a term $a_{\mathbf{i}} X^{\mathbf{i}}$ as $\text{val}_{\mathbf{r}}(a_{\mathbf{i}} X^{\mathbf{i}}) = \text{val}(a_{\mathbf{i}}) - \mathbf{r} \cdot \mathbf{i}$, and the Gauss valuation of $\sum a_{\mathbf{i}} X^{\mathbf{i}} \in K\{X; \mathbf{r}\}$ as the minimum of the Gauss valuations of its terms. The valuation defines a metric on $K\{X; \mathbf{r}\}$, for which a sequence $(f_n)_{n \in \mathbb{N}} \in K\{X; \mathbf{r}\}$ converges to zero iff $\text{val}_{\mathbf{r}}(f_n) \xrightarrow{n \rightarrow +\infty} +\infty$.

In this article, we shall frequently need to consider all terms with minimal valuation together.

Definition 2.1. Let $f \in \sum_{\alpha \in \mathbb{N}^n} c_{\alpha} X^{\alpha} \in K\{X; \mathbf{r}\}$. The \mathbf{r} -support of f is

$$\text{Supp}_{\mathbf{r}}(f) = \{\alpha \text{ s.t. } \text{val}_{\mathbf{r}}(c_{\alpha} X^{\alpha}) = \text{val}_{\mathbf{r}}(f)\},$$

and the *initial part* of f is

$$\text{in}_{\mathbf{r}}(f) = \sum_{\alpha \in \text{Supp}_{\mathbf{r}}(f)} c_{\alpha} X^{\alpha}.$$

By definition, for $f \in K\{X; \mathbf{r}\}$, $\text{in}_{\mathbf{r}}(f)$ is a polynomial.

We fix a classical *monomial order* \leq_m on the set of monomials $X^{\mathbf{i}}$, which will be used for tie-breaks. Given two terms $aX^{\mathbf{i}}$ and $bX^{\mathbf{j}}$ (with $a, b \in K^{\times}$), we write $aX^{\mathbf{i}} <_{\mathbf{r},m} bX^{\mathbf{j}}$ if $\text{val}_{\mathbf{r}}(aX^{\mathbf{i}}) > \text{val}_{\mathbf{r}}(bX^{\mathbf{j}})$, or $\text{val}_{\mathbf{r}}(aX^{\mathbf{i}}) = \text{val}_{\mathbf{r}}(bX^{\mathbf{j}})$ and $X^{\mathbf{i}} <_m X^{\mathbf{j}}$. By definition, the leading term of a Tate series $f = \sum a_{\mathbf{i}} X^{\mathbf{i}} \in K\{X; \mathbf{r}\}$ is its maximal term, and is denoted by $\text{LT}_{\mathbf{r},m}(f)$. Its coefficient and its monomial are denoted $\text{LC}_{\mathbf{r},m}(f)$ and $\text{LM}_{\mathbf{r},m}(f)$, with $\text{LT}_{\mathbf{r},m}(f) = \text{LC}_{\mathbf{r},m}(f) \times \text{LM}_{\mathbf{r},m}(f)$. For $f, g \in K\{X; \mathbf{r}\}$, we define their S -polynomial as

$$\text{S-Poly}(f, g) = \frac{\text{LT}_{\mathbf{r},m}(g)}{D(f, g)} f - \frac{\text{LT}_{\mathbf{r},m}(f)}{D(f, g)} g$$

where $D(f, g) = \text{gcd}(\text{LT}_{\mathbf{r},m}(f), \text{LT}_{\mathbf{r},m}(g))$.

A Gröbner basis (or GB for short) of an ideal I of $K\{X; \mathbf{r}\}$ is a set $G \subseteq I$ such that for all $f \in I$, there exists an index $g \in G$ such that $\text{LT}_{\mathbf{r},m}(g)$ divides $\text{LT}_{\mathbf{r},m}(f)$. A finite Gröbner basis (g_1, \dots, g_s)

is *reduced* if all $\text{LT}_{\mathbf{r},m}(g_i)$'s are monic, minimally generate $\text{LT}_{\mathbf{r},m}(I)$ and, for any i , $\text{LT}_{\mathbf{r},m}(g_i)$ is the only term of g_i in $\text{LT}_{\mathbf{r},m}(I)$.

The following theorem was proved in [4].

Theorem 2.2. *Let I be an ideal of $K\{X; \mathbf{r}\}$, then I admits a finite Gröbner basis.*

2.2 Local Gröbner bases of polynomial ideals in Tate algebras

For a polynomial ideal $I \subset K[X]$ and a system of convergence log-radii \mathbf{r} , we define $I_{\mathbf{r}}$ to be the ideal of $K\{X; \mathbf{r}\}$ generated by the polynomials of I . It is the completion of I with respect to $\text{val}_{\mathbf{r}}$. The ideal $I_{\mathbf{r}}$ usually contains many series and polynomials not in I . However, as I is dense in $I_{\mathbf{r}}$, it was proved in [7] that $I_{\mathbf{r}}$ does not contain more leading terms than I , and that $I_{\mathbf{r}}$ admits a Gröbner basis comprised of polynomials.

Definition 2.3. Let $I \subset K[X]$ be an ideal, \mathbf{r} a system of log-radii and $I_{\mathbf{r}}$ the completion of I in $K\{X; \mathbf{r}\}$. An \mathbf{r} -local Gröbner basis of I is a Gröbner basis of $I_{\mathbf{r}}$ comprised only of polynomials.

If one needs to vary the convergence log-radii, the following object is of interest:

Definition 2.4. Let $I \subset K[X]$ be an ideal. A finite set $G \subset I \subset K[X]$ such that for any $\mathbf{r} \in \mathbb{Q}^n$, G is an \mathbf{r} -local GB of I is called a *universal analytic Gröbner basis* of I (UAGB for short).

In the usual setting, it is required that universal Gröbner bases be a Gröbner basis for all monomial orders. Here, the definition requires only that all convergence radii be covered, without any restriction on the tie-breaking monomial order. However, we prove in Lemma 3.4 that given a polynomial ideal, any term ordering can be achieved with a suitable choice of a system of convergence radii. Finally [7] culminated with the following result:

Theorem 2.5. *Let $I \subset K[X]$ be an ideal. Then there exists a universal analytic Gröbner basis of I .*

While the proof was not constructive, we provide in this article an algorithm to compute a UAGB of any polynomial ideal.

2.3 Homogenization and dehomogenization

Our algorithm to compute a UAGB of a polynomial ideal will rely on homogenization and dehomogenization. We consign here notations and basic properties taken from [7, §3.3]

Definition 2.6. Let $(\cdot)^*$ and $(\cdot)_*$ be the homogenization and dehomogenization applications between $K[X]$ and $K[X, t]$. If $I \subset K[X]$ is an ideal, we define $I^* \subset K[X, t]$ to be the ideal spanned by the f^* for $f \in I$.

Given $\mathbf{r} \in \mathbb{Q}^n$ and \leq_m a monomial order, we extend the term order $<_{\mathbf{r},m}$ to $K[X, t]$ and $K\{X, t; \mathbf{r}, 0\}$ as follows.

Definition 2.7. Given two terms $aX^{\alpha}t^u$ and $bX^{\beta}t^v$, we write that $aX^{\alpha}t^u <_{(\mathbf{r},0),m} bX^{\beta}t^v$ if one of the following holds:

- $\text{val}_{\mathbf{r}}(aX^{\alpha}) > \text{val}_{\mathbf{r}}(bX^{\beta})$ (i.e., $\text{val}_{\mathbf{r},0}(aX^{\alpha}t^u) > \text{val}_{\mathbf{r},0}(bX^{\beta}t^v)$).
- $\text{val}_{\mathbf{r}}(aX^{\alpha}) = \text{val}_{\mathbf{r}}(bX^{\beta})$ and $\deg(X^{\alpha}t^u) < \deg(X^{\beta}t^v)$.
- $\text{val}_{\mathbf{r}}(aX^{\alpha}) = \text{val}_{\mathbf{r}}(bX^{\beta})$, $\deg(X^{\alpha}t^u) = \deg(X^{\beta}t^v)$ and $X^{\alpha} <_m X^{\beta}$.

This defines a term order on $K\{X, t; \mathbf{r}, 0\}$.

With this order, dehomogenization preserves leading terms of homogeneous polynomials of $K[X, t]$.

Lemma 2.8 (Lem. 3.5 in [7]). *Let $\mathbf{r} \in \mathbb{Q}^n$. Let $h \in K[X, t]$ be a homogeneous polynomial. Then $\text{LT}_{(\mathbf{r}, 0), m}(h)_* = \text{LT}_{\mathbf{r}, m}(h_*)$. Let $f \in K[X]$, then $\text{LT}_{\mathbf{r}, m}(f) = (\text{LT}_{(\mathbf{r}, 0), m}(f^*))_*$.*

3 TERM ORDERS

3.1 Convergence radii and term orders

In this section, we collect different results regarding term orders in Tate algebras. First, we consider the relation between term orders and systems of convergence log-radii, and show that given finite data (e.g. a finite set of polynomials or an ideal), it is always possible to realize a term order by a suitable choice of system of convergence log-radii.

Definition 3.1. Let $F = (f_1, \dots, f_s) \in K[X]^s$. Given a term order $<$, we define

$$\text{LT}_<(F) = \{\text{LT}_<(f) : f \in F\}.$$

We say that two term orders $<_1$ and $<_2$ on $K[X]$ are equivalent with respect to F if

$$\text{LT}_{<_1}(F) = \text{LT}_{<_2}(F).$$

Let $I \subseteq K[X]^s$ be an ideal, we say that two term orders $<_1$ and $<_2$ on $K[X]$ are equivalent with respect to I if

$$\{\text{LT}_{<_1}(f) : f \in I\} = \{\text{LT}_{<_2}(f) : f \in I\}$$

or equivalently

$$\text{LT}_{<_1}(I) = \text{LT}_{<_2}(I).$$

The two relations are connected as follows.

Lemma 3.2. *Let $I \subseteq K[X]^s$, and $<_1, <_2$ be two term orders. Let G be a Gröbner basis of I w.r.t. $<_1$ and $<_2$. If $<_1$ and $<_2$ are equivalent w.r.t. G , then they are equivalent w.r.t. I .*

PROOF. Let $<_1$ and $<_2$ be two term orders equivalent w.r.t. G . Since G is a GB of I w.r.t. $<_1$ and $<_2$, $\text{LT}_{<_1}(I)$ is spanned by $\text{LT}_{<_1}(G)$, which is equal to $\text{LT}_{<_2}(G)$ spanning $\text{LT}_{<_2}(I)$. \square

The following lemma states that modulo equivalence w.r.t. F , it is always possible to choose a term order determined only by the convergence condition.

Lemma 3.3. *Let $\mathbf{r} \in \mathbb{Q}^n$ and let $\leq_{\mathbf{r}, m}$ be a term order defined by $\text{val}_{\mathbf{r}}$ and a tie-breaking order \leq_m . Let T be a finite set of terms in $K[X]$. There is an $\mathbf{s} \in \mathbb{Q}^n$ such that for all $t_1, t_2 \in T$,*

$$t_1 > t_2 \iff \text{val}_{\mathbf{s}}(t_1) < \text{val}_{\mathbf{s}}(t_2).$$

In particular, if F is a finite set in $K[X]$, any equivalence class of term orders w.r.t. F contains a term order $<_{\mathbf{s}, m}$ such that for all $f \in F$, $\text{LT}_{\mathbf{s}, m}(f) = \text{in}_{\mathbf{s}}(f)$.

PROOF. Thanks to [20, Th. 1] (see also [12, Lem. 1.3.1]), there exists some $\mathbf{u} \in \mathbb{Q}^n$ such that for all $c_1 X^\alpha \neq c_2 X^\beta$ in T ,

$$X^\alpha >_m X^\beta \iff \alpha \cdot \mathbf{u} > \beta \cdot \mathbf{u}.$$

By considering the finite set of pairs of terms in T , there is a small enough $\varepsilon \in \mathbb{Q}_{>0}$ such that for any t_1, t_2 in T , if $\text{val}_{\mathbf{r}}(t_1) < \text{val}_{\mathbf{r}}(t_2)$ then $\text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_1) < \text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_2)$.

Let $\mathbf{s} = \mathbf{r} - \varepsilon\mathbf{u}$. Then, for any $t_1 = c_1 X^\alpha, t_2 = c_2 X^\beta$ in T such that $t_1 >_{\mathbf{r}, m} t_2$, one of the following is true:

- $\text{val}_{\mathbf{r}}(t_1) < \text{val}_{\mathbf{r}}(t_2)$ and therefore $\text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_1) < \text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_2)$;
- $\text{val}_{\mathbf{r}}(t_1) = \text{val}_{\mathbf{r}}(t_2)$ and $X^\alpha >_m X^\beta$, but then $\alpha \cdot \mathbf{u} > \beta \cdot \mathbf{u}$ and $\text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_1) < \text{val}_{\mathbf{r}-\varepsilon\mathbf{u}}(t_2)$.

Finally, if $\alpha = \beta$ and $\text{val}(c_1) = \text{val}(c_2)$, then $c_1 X^\alpha \not> c_2 X^\beta$ and there is nothing to prove. Therefore, \mathbf{s} satisfies the claim.

The consequence for equivalence classes w.r.t. F follows, by setting $T = \bigcup_{f \in F} \text{Supp}(f)$. \square

Lemma 3.4. *Let $I \subseteq K[X]$ be an ideal, $<_{\mathbf{r}, m_1}$ be a term order and G a local Gröbner basis of I w.r.t. $<_{\mathbf{r}, m_1}$. Then there exists $\mathbf{s} \in \mathbb{Q}^n$ such that, for any tie-break order $<_{m_2}$:*

- G is a Gröbner basis of I w.r.t. $<_{\mathbf{s}, m_2}$;
- $<_{\mathbf{r}, m_1}$ is equivalent to $<_{\mathbf{s}, m_2}$ w.r.t. I ;
- $\text{LT}_{\mathbf{s}, m_2}(I) = \langle \text{in}_{\mathbf{s}}(f) : f \in I \rangle$.

PROOF. Let T be the set of all terms which appear in G and in the course of Buchberger's algorithm with weak normal form with G as input w.r.t. the $<_{\mathbf{r}, m_1}$ ordering. By Lemma 3.3, there exists \mathbf{s} such that all terms in T have distinct \mathbf{s} -valuation, compatible with the order $<_{\mathbf{r}, m_1}$. Let $<_{m_2}$ be a tie-break order.

Note that when running Buchberger's algorithm with WNF, the radii of convergence are only used for determining leading terms. So if we run the algorithm with G as input and for the order $<_{\mathbf{s}, m_2}$, all comparisons will be the same, the exact same terms will appear, and the result will be the same: all the S-polynomials have a weak normal form of 0 w.r.t. G , and so G is a GB of I w.r.t. $<_{\mathbf{s}, m_2}$.

By Lemma 3.2, this, together with the fact that the elements of G have the same leading term for both orders, implies that the term orders are equivalent w.r.t. I .

Clearly, since for all $g \in G$, $\text{LT}_{\mathbf{s}, m_2}(g) = \text{in}_{\mathbf{s}}(g)$, $\text{LT}_{\mathbf{s}, m_2}(I) \subseteq \langle \text{in}_{\mathbf{s}}(f) : f \in I \rangle$. For the reverse inclusion, let $f \in I$, and write $\text{in}_{\mathbf{s}}(f) = \text{LT}_{\mathbf{s}, m_2}(f) + r$. Since G is a GB of I w.r.t. $<_{\mathbf{s}, m_2}$, there exists a $g \in G$ and a term τ such that $\text{LT}_{\mathbf{s}, m_2}(f) = \tau \text{LT}_{\mathbf{s}, m_2}(g)$. Since all other terms in g have strictly higher valuation, then either $\text{val}_{\mathbf{s}}(f - \tau g) > \text{val}_{\mathbf{s}}(f)$ and then $r = 0$, or $\text{val}_{\mathbf{s}}(f - \tau g) = \text{val}_{\mathbf{s}}(f)$ and $\text{in}_{\mathbf{s}}(f - \tau g) = r = \text{in}_{\mathbf{s}}(f) - \text{LT}_{\mathbf{s}, m_2}(f)$. Repeating the process until $r = 0$, we get that $\text{in}_{\mathbf{s}}(f)$ is a linear combination of terms divisible by $\text{LT}_{\mathbf{s}, m_2}(g)$ for some g 's, and therefore $\text{LT}_{\mathbf{s}, m_2}(I) = \langle \text{in}_{\mathbf{s}}(f) : f \in I \rangle$. \square

For such an \mathbf{s} , we say that \mathbf{s} defines a term order for I . The tie-breaking order $<_{m_2}$ becomes irrelevant, and we omit it from the notations. In the rest of the paper, unless specified otherwise, we will always be considering orders $<_{\mathbf{s}}$ where \mathbf{s} defines a term order.

3.2 Newton polytopes

In this section, we generalize known results from the classical case to orderings compatible with the valuation, and relating equivalence classes of term orderings w.r.t. a finite set of polynomials, with data from its Newton polytope. Let $F = (f_1, \dots, f_s) \in K[X]^s$.

Definition 3.5. For $f \in K[X]$, we define $\mathcal{N}(f)$ to be the convex hull of

$$\{(\text{val}(c)), \alpha_1, \dots, \alpha_n \text{ for } cX^\alpha \in \text{Supp}(f)\} + \mathbb{R}_+(1, 0, \dots, 0).$$

We then define $\mathcal{N}(F)$ to be the Minkowski sum of the $\mathcal{N}(f_i)$'s.

Remark 3.6. For $n = s = 1$, this coincides with the classical definition of the Newton polygon (up to a symmetry).

Lemma 3.7. *For the convex polyhedron $\mathcal{N}(F)$, $\beta \in \mathcal{N}(F)$ is a vertex if and only if there is some $U = (1, u_1, \dots, u_n) \in \mathbb{Q}^{n+1}$ such that $\beta \cdot U$ is the unique minimum of the $\alpha \cdot U$'s for $\alpha \in \mathcal{N}(F)$.*

PROOF. Since $\mathcal{N}(F)$ is a convex polyhedron, β is a vertex if and only if there is some $U = (u_0, u_1, \dots, u_n) \in \mathbb{Q}^{n+1}$ such that $\beta \cdot U$ is the unique minimum of the $\alpha \cdot U$'s for $\alpha \in \mathcal{N}(F)$. Since $\mathcal{N}(F)$ is defined from half-lines of the form $(\text{val}(c_{i,k}), \alpha_{i,k}^{(1)}, \dots, \alpha_{i,k}^{(n)}) + \mathbb{R}_+(1, 0, \dots, 0)$, then we can further assume that the U in the previous equivalence is such that $u_0 > 0$ and then by multiplying by a positive rational, we can assume that $U = (1, u_1, \dots, u_n)$. \square

Proposition 3.8. *The vertices of $\mathcal{N}(F)$ are in one-to-one correspondence with the equivalence classes of term orders with respect to F .*

PROOF. For any $i \in [1, s]$, we write $f_i = \sum_{j=1}^{l_i} c_{i,j} X^{\alpha_{i,j}}$. Let us define the following set of index vectors:

$$J := \{(j_1, \dots, j_s) \in \mathbb{N}^s : \forall i \in [1, s], 1 \leq j_i \leq l_i\}.$$

For $\mathbf{j} \in J$, and $i \in [1, s]$, we define $D_{i,j_i} = [1, l_i] \setminus \{j_i\}$ and we define

$$C_j = \{\mathbf{r} \in \mathbb{Q}^n : \forall i \in [1, s], \forall j \in D_{i,j_i}, \alpha_{i,j_i} \cdot (1, \mathbf{r}) < \alpha_{i,j} \cdot (1, \mathbf{r})\},$$

so that for any $\mathbf{r} \in C_j$ and $i \in [1, s]$, $\text{LT}_{\mathbf{r}}(f_i) = c_{i,j_i} X^{\alpha_{i,j_i}}$. Then, thanks to Lemma 3.3, there is one equivalence class of term orders with respect to F for each non empty C_j .

Being defined as a Minkowski sum, $\mathcal{N}(F)$ is the convex hull of the rays $\alpha_j + \mathbb{R}_+(1, 0, \dots, 0)$ for $\alpha_j := \sum_{i=1}^s \alpha_{i,j_i}$ and $\mathbf{j} \in J$. Thus, its vertices are among the α_j 's. Thanks to Lemma 3.7, α_j is a vertex of $\mathcal{N}(F)$ if and only if there is some $U = (1, u_1, \dots, u_n) \in \mathbb{Q}^{n+1}$ such that $\alpha_j \cdot U$ is the unique minimum of the $\alpha \cdot U$'s for $\alpha \in \mathcal{N}(F)$. Consequently, if $\mathbf{j}' = \mathbf{j} + (0, \dots, 0, j'_i, 0, \dots, 0) - (0, \dots, 0, j_i, 0, \dots, 0)$ only differs from \mathbf{j} on the coordinate i (for some $j'_i \in D_{i,j_i}$), then we can deduce from $\alpha_j \cdot U < \alpha_{j'} \cdot U$ that $\alpha_{i,j_i} \cdot U < \alpha_{i,j'_i} \cdot U$. It implies that $(u_1, \dots, u_n) \in C_j$. And the converse is true: if $(u_1, \dots, u_n) \in C_j$ then α_j is a vertex of $\mathcal{N}(F)$.

This is enough to conclude that there is a one-to-one correspondence between vertices of $\mathcal{N}(F)$ and equivalence classes of term orders with respect to F . \square

4 UNIVERSAL ANALYTIC GRÖBNER BASES¹

4.1 Testing whether a set is a UAGB

The results of Section 3.2 are enough to immediately provide us with a procedure for deciding whether a set is a UAGB (Algorithm 1).

Proposition 4.1. *$F = (f_1, \dots, f_s)$ is a UAGB of $I = \langle F \rangle$ if and only if for any \mathbf{r} in the equivalence classes of term orders with respect to F , F is a GB of $I_{\mathbf{r}}$. In particular, Algorithm 1 is correct.*

4.2 Computing a UAGB

We now show how to use that procedure to compute a UAGB. To that end, we recall the following result from [7].

Theorem 4.2. [7, Thm 7.6] *Let $I \subset K[X]$ be an ideal. Then the set $\text{Terms}(I) := \{\text{LT}(I_{\mathbf{r}}) \text{ for } \mathbf{r} \in \mathbb{Q}^n\}$ is finite.*

¹A toy implementation of the algorithms in this section is available at: <https://gist.github.com/TristanVaccon>

Algorithm 1 TestUAGB

Input: $F = \{f_1, \dots, f_s\} \in K[X]$ generating $I \subseteq K[X]$

Output: True if F is a UAGB of I , otherwise (False, \mathbf{u}) with $\mathbf{u} \in \mathbb{Q}^n$ such that F is not a \mathbf{u} -local GB of I

```

1: Compute  $N = \mathcal{N}(f_1, \dots, f_s)$ ;
2: for  $\beta \in \{\text{vertices of } N\}$  do
3:   Compute  $U = (1, \mathbf{u})$  characterizing  $\beta$  as in Lemma 3.7 ;
4:   Compute  $G_{\mathbf{u}}$ , a  $\mathbf{u}$ -local GB of  $I$  ;
5:   if  $\exists g \in G_{\mathbf{u}}$  not reducible modulo  $F$  for the order  $<_{\mathbf{u}}$  then
6:     return (False,  $\mathbf{u}$ ) ;
7: return True ;
```

The proof of the previous theorem relies on the following lemma, which we also need.

Lemma 4.3. *Let $I \subset K[X]$ be an ideal. Let \leq_1 and \leq_2 be two term orders such that $\text{LT}_{\leq_1}(I) = \text{LT}_{\leq_2}(I)$. Let $G \subset I$ be a reduced (local) GB of I w.r.t. \leq_1 . Then G is also a reduced GB of I w.r.t. \leq_2 .*

PROOF. Let $g \in G$. Since G is reduced and $\text{LT}_{\leq_1}(I) = \text{LT}_{\leq_2}(I)$, then the only term of g belonging to $\text{LT}_{\leq_2}(I)$ is $\text{LT}_{\leq_1}(g)$. Thus $\text{LT}_{\leq_1}(g) = \text{LT}_{\leq_2}(g)$, and G is a reduced GB of I w.r.t. \leq_2 . \square

Unlike in the classical case, it is not in general possible to guarantee that any polynomial ideal admits a reduced local Gröbner basis for any convergence radii. However, for homogeneous ideals, [7, Lem. 7.2] guarantees that there is a reduced local GB comprised only of polynomials, which can then be computed using any GB algorithm from [4, 5, 7, 8, 26–28].

The proofs in [7] relied on homogenization and dehomogenization of ideals and were not constructive. In this paper, we replace computations in the homogenized ideal by computations in the ideal spanned by the homogenization of its generators.

Notation 4.4. Let $F = (f_1, \dots, f_s) \in K[X]^s$, we define

$$F^h = (f_1^*, \dots, f_s^*).$$

In general, $\langle F^h \rangle \subseteq I^*$ but $I = \langle F^h \rangle_* = (I^*)_*$. By [7, Lem 7.5], the dehomogenization of a GB of $(I^*)_{(\mathbf{r},0)}$ made of homogeneous polynomials of $K[X, t]$ is a polynomial GB of $I_{\mathbf{r}}$. This result is still true for $\langle F^h \rangle$, by the following lemma.

Lemma 4.5. *Let $F = (f_1, \dots, f_s) \in K[X]^s$, $I = \langle F \rangle$ and $\mathbf{r} \in \mathbb{Q}^n$. Let (h_1, \dots, h_s) be a finite Gröbner basis of $\langle F^h \rangle_{(\mathbf{r},0)} \subset K[X, t; \mathbf{r}, 0]$ made of homogeneous polynomials of $\langle F^h \rangle$ (hence in $K[X, t]$). Then $(h_{1,*}, \dots, h_{s,*})$ is an \mathbf{r} -local GB of I .*

PROOF. Firstly, due to being a dehomogenization of homogeneous elements of $\langle F^h \rangle$, the $h_{i,*}$'s are in I (it is enough to dehomogenize an homogeneous combination of the f_i^* 's).

Secondly, by [7, Cor 5.4], it is enough to check that for any $f \in I$, $\text{LT}_{\mathbf{r}}(f)$ is divisible by one of the $\text{LT}_{\mathbf{r}}(h_{i,*})$'s.

Let $f \in I$. Since $I = \langle F^h \rangle_*$, there is some homogeneous polynomial $g \in \langle F^h \rangle$ such that $g_* = f$. Then $g \in \langle F^h \rangle \subset (\langle F^h \rangle)_{(\mathbf{r},0)}$ so there is some i such that $\text{LT}_{(\mathbf{r},0)}(h_i)$ divides $\text{LT}_{(\mathbf{r},0)}(g)$. Then thanks to Lemma 2.8, $\text{LT}_{\mathbf{r}}(f) = \text{LT}_{(\mathbf{r},0)}(g)_*$, $\text{LT}_{\mathbf{r}}(h_{i,*}) = \text{LT}_{(\mathbf{r},0)}(h_i)_*$, and monomial divisibility is preserved by dehomogenization. So $\text{LT}_{\mathbf{r}}(h_{i,*})$ divides $\text{LT}_{\mathbf{r}}(f)$ and the proof is complete. \square

Remark 4.6. One needs to be careful that the dehomogenization of a *reduced* Gröbner basis of $\langle F^h \rangle_{(r,0)}$ (made of homogeneous polynomials of $\langle F^h \rangle$) need not even be *minimal*, one leading monomial may divide another after dehomogenization (e.g. yt^3 and x^2yt).

We can now provide an algorithm for computing UAGBs.

Algorithm 2 UAGB

Input: $F \in K[X]^s$, generating $I \in K[X]$.

Output: G a UAGB for I .

```

1:  $G := \{f^* \text{ for } f \in F\}$  (and define  $J = \langle F^h \rangle$ );
2: while TestUAGB( $G$ ) is not True do
3:    $\mathbf{r} :=$  system of log radii such that  $G$  is not a GB of  $J_{\mathbf{r}}$  (as
     produced by TestUAGB);
4:    $H :=$  ReducedGB( $G, \mathbf{r}$ ); // Polynomial reduced local GB
5:    $G := G \cup H$ ;
6: return  $\{g^* \text{ for } g \in G\}$ ;

```

Theorem 4.7. *Algorithm 2 is correct and computes a UAGB in finite time. Furthermore, if the input polynomials are homogeneous, the UAGB contains a reduced Gröbner basis for all orders.*

PROOF. Thanks to Theorem 4.2, if $J = \langle F^h \rangle$, there is an integer t and a finite set of term orders \leq_1, \dots, \leq_t such that $\text{Terms}(J)$ is given by $\{\text{LT}(J_{\leq_1}), \dots, \text{LT}(J_{\leq_t})\}$. Let us assume that G contains a reduced GB w.r.t. each of the orders \leq_1, \dots, \leq_l for some $l \leq t$.

Let us assume that TestUAGB(G) fails because G is not an \mathbf{r} -local GB of I for some \mathbf{r} . Then, thanks to Theorem 4.2 and Lemma 4.3, $\text{LT}(J_{\leq_i}) \neq \text{LT}(J_{\mathbf{r}})$ for any integer i , $1 \leq i \leq l$ and $\text{LT}(J_{\leq_j}) = \text{LT}(J_{\mathbf{r}})$ for some integer j , $l < j \leq t$. Up to renumbering, we may assume that $j = l + 1$. Let H be the reduced GB of J for \mathbf{r} . Then $G \cup H$ contains a reduced GB for the orders $\leq_1, \dots, \leq_{l+1}$.

We then prove by induction that, after at most t calls to ReducedGB and to TestUAGB, the algorithm outputs G such that G contains a reduced GB of J for each of \leq_1, \dots, \leq_t and hence, is a UAGB of J .

Finally, thanks to Lemma 4.5, the dehomogenization of G is a UAGB of $I = J_*$. If the input polynomials are homogeneous, the homogenization and dehomogenization steps are trivial, and the property that the UAGB contains a reduced GB for all orders is preserved. \square

Remark 4.8. From the proof of Theorem 4.7, Algorithm 2 needs at most $\#\text{Terms}(J)$ loops to compute a UAGB. Each loop may however cause many GB computations as it is unclear how the edges of the Newton polytopes vary along the computation.

4.3 Examples

Example 4.9. Let $F = [x - 7y, y - 7y^2]$ in $\mathbb{Q}_7[x, y]$. Then in $\mathbb{Q}_7[x, y, t]$, one finds that F^h is not a GB for the weight $[0, 2, 0]$. The corresponding reduced GB will add the polynomial $x^2 - xt$ and $G_h = [x - 7y, yt - 7y^2, x^2 - xt]$ is then a UAGB for $\langle F^h \rangle$. Thus, $G = [x - 7y, y - 7y^2, x^2 - x]$ is a UAGB of $\langle F \rangle$.

Remark 4.10. No finite approximate interreduction \overline{F} of $[x - 7y, y - 7y^2]$ is enough to be a UAGB.

5 TROPICAL GEOMETRY

5.1 Analytical tropical varieties

In this section, we show that tropical geometry on Tate polynomial ideals specializes that of classical polynomial ideals. In particular, the results of Section 3 give us a Tate analogue of the Gröbner fan, and allow us to generalize the results of [17, 22] for computing tropical varieties in $K[[T]][X]$, to any Tate algebra.

First, we recall the classical notions of tropical geometry, and state their natural generalization to Tate algebras.

Definition 5.1. Let $\mathbf{w} = (w_0, \dots, w_n) \in \mathbb{R}_{<0} \times \mathbb{R}^n$ be a system of weights. For a monomial $m = X_1^{\alpha_1} \dots X_n^{\alpha_n}$ and $c \in K$, we define its weighted degree

$$\deg_{\mathbf{w}}(cX_1^{\alpha_1} \dots X_n^{\alpha_n}) = w_0 \text{val}(c) + \sum_{i=1}^n w_i \alpha_i.$$

For $f \in K[X]$, let $\deg_{\mathbf{w}}(f) = \max(\deg_{\mathbf{w}}(t) : t \in \text{Supp}(f))$. We define the initial form of f as

$$\text{in}_{\mathbf{w}}(f) = \sum \{t : t \in \text{Supp}(f), \deg_{\mathbf{w}}(t) = \deg_{\mathbf{w}}(f)\}.$$

Let $I \subset K[X]$ be an ideal. Then $\text{in}_{\mathbf{w}}(I)$, the initial ideal of I with respect to a system of weights \mathbf{w} , is the ideal spanned by all $\text{in}_{\mathbf{w}}(f)$ for f in I . The *tropical variety* associated to I is then defined as

$\text{trop}(I) = \{\mathbf{w} \in \mathbb{R}_{<0} \times \mathbb{R}^n : \text{in}_{\mathbf{w}}(I) \text{ does not contain a monomial}\}.$

We say that the system of weights \mathbf{w} and the system of log-radii \mathbf{r} are *compatible* if $\mathbf{r} = -\left(\frac{w_1}{w_0}, \dots, \frac{w_n}{w_0}\right)$. Conversely, given a system of log-radii \mathbf{r} , the system of weights $(-1, r_1, \dots, r_n)$ is compatible with \mathbf{r} . The definitions above extend naturally to series and ideals in $K\{X; \mathbf{r}\}$ by restricting to systems of weights which are compatible with \mathbf{r} .

Remark 5.2. In particular, $\text{trop}(I_{\mathbf{r}})$ is either empty or a half-line formed of all the systems of weights compatible with \mathbf{r} .

If the systems of weights \mathbf{w} and the system of log-radii \mathbf{r} are compatible, then for any term t ,

$$\deg_{\mathbf{w}}(t) = w_0 \text{val}_{\mathbf{r}}(t).$$

This implies that $\deg_{\mathbf{w}}$ is a graduation: for any terms t, t' , $\deg_{\mathbf{w}}(t + t') \leq \max(\deg_{\mathbf{w}}(t), \deg_{\mathbf{w}}(t'))$ with equality if $\deg_{\mathbf{w}}(t) \neq \deg_{\mathbf{w}}(t')$, and $\deg_{\mathbf{w}}(tt') = \deg_{\mathbf{w}}(t) + \deg_{\mathbf{w}}(t')$.

The main result of this section is the fact that the tropical variety associated to I is the *union* of the tropical varieties of all its completions $I_{\mathbf{r}}$.

Lemma 5.3. *Let \mathbf{w} be a system of weights, let \mathbf{r} be the compatible system of convergence radii and let \leq_m be a monomial order. Let $f \in K\{X; \mathbf{r}\}$. Then:*

- (1) $\text{in}_{\mathbf{w}}(f) = \text{in}_{\mathbf{r}}(f)$, and in particular it is a polynomial;
- (2) $\text{LT}_{\leq_m}(\text{in}_{\mathbf{w}}(f)) = \text{LT}_{\mathbf{r}, \leq_m}(f)$;

PROOF. By compatibility between the system of weights and the convergence log-radii, $\text{val}_{\mathbf{r}}(aX^i) = w_0^{-1} \deg_{\mathbf{w}}(aX^i)$, and $\text{val}_{\mathbf{r}}(f) = w_0^{-1} \deg_{\mathbf{w}}(f)$. So $\text{in}_{\mathbf{w}}(f)$ is the sum of all terms with minimal Gauss valuation in the support of f , which is by definition $\text{in}_{\mathbf{r}}(f)$. The rest follows from the convergence properties in Tate series and the definition of the term order. \square

Theorem 5.4. Let $I \subseteq K[X]$ be an ideal. Let $\mathbf{w} \in \mathbb{R}_{<0} \times \mathbb{R}^n$ be a system of weights, and let $\mathbf{r} = -(\mathbf{w}_1/\mathbf{w}_0, \dots, \mathbf{w}_n/\mathbf{w}_0) \in \mathbb{R}^n$ be the compatible system of convergence log-radii. Let $I_{\mathbf{r}} \subseteq K\{X; \mathbf{r}\}$ be the completion of I . Then

$$\text{in}_{\mathbf{w}}(I) = \text{in}_{\mathbf{w}}(I_{\mathbf{r}}) \cap K[X]$$

and in particular, $\mathbf{w} \in \text{trop}(I)$ if and only if $\mathbf{w} \in \text{trop}(I_{\mathbf{r}})$. Globally,

$$\text{trop}(I) = \bigcup_{s \in \mathbb{R}^n} \text{trop}(I_s).$$

PROOF. Clearly $\text{in}_{\mathbf{w}}(I) \subseteq \text{in}_{\mathbf{w}}(I_{\mathbf{r}}) \cap K[X]$. Conversely, let us first consider $f \in I_{\mathbf{r}}$ and let f_1, \dots, f_k be polynomials generating I .

There exist series g_1, \dots, g_k such that $f = g_1 f_1 + \dots + g_k f_k$. Let $d = \text{val}_{\mathbf{r}}(f)$, and write each series g_i as $h_i + r_i$, where h_i is the sum of all terms with Gauss valuation at most $d - \text{val}_{\mathbf{r}}(f_i)$ and $r_i = g_i - h_i$ has Gauss valuation greater than $d - \text{val}_{\mathbf{r}}(f_i)$. By the convergence property, the h_i 's are polynomials. The decomposition of f becomes

$$f = h_1 f_1 + \dots + h_k f_k + (r_1 f_1 + \dots + r_k f_k)$$

where the latter group consists exclusively of terms with Gauss valuation greater than d . So none of those terms can appear in the initial form of f , and as a consequence, $\text{in}_{\mathbf{w}}(f) = \text{in}_{\mathbf{w}}(h_1 f_1 + \dots + h_k f_k)$. Since the h_i are polynomials, $h_1 f_1 + \dots + h_k f_k \in I$, and therefore $\text{in}_{\mathbf{w}}(f) \in \text{in}_{\mathbf{w}}(I)$.

Now let $h \in \text{in}_{\mathbf{w}}(I) \cap K[X]$, there exists series $g_1, \dots, g_l \in I_{\mathbf{r}}$ and series q_1, \dots, q_l such that

$$h = q_1 \text{in}_{\mathbf{w}}(g_1) + \dots + q_l \text{in}_{\mathbf{w}}(g_l).$$

From the above, we know that $\text{in}_{\mathbf{w}}(g_i) \in \text{in}_{\mathbf{w}}(I)$ for all i . Since h is a polynomial, it has a maximal Gauss valuation d . Similarly to before, any term in q_i with Gauss valuation greater than $d - \text{val}_{\mathbf{r}}(g_i)$ cannot appear in h , so those terms must add to zero on the right hand side, and we can assume that the cofactors q_i are polynomials, and therefore $h \in \text{in}_{\mathbf{w}}(I)$ and the second inclusion is proved.

The rest of the statement follows by definition. \square

5.2 Analytic Gröbner fan

Similarly to the polynomial case, tropical varieties can be computed using the Gröbner fan of the ideal. In this section, we recall those definitions.

The relation between tropical varieties and Gröbner fans is the same as in the usual case, namely, that initial forms generalize leading terms.

Definition 5.5. Let $\mathbf{w} \in \mathbb{R}_{<0} \times \mathbb{R}^n$ be a system of weights. Let \leq be a term order on $K[X]$. We say that \leq refines \mathbf{w} if for all terms t_1, t_2 , $\deg_{\mathbf{w}}(t_1) \geq \deg_{\mathbf{w}}(t_2) \implies t_1 \geq t_2$. We say that \mathbf{w} defines a term order for a finite set of polynomials or an ideal if $\mathbf{r} = -(\mathbf{w}_1/\mathbf{w}_0, \dots, \mathbf{w}_n/\mathbf{w}_0)$ defines a term order for that set or ideal.

If \mathbf{w} defines a term order for a finite set F , for all $f \in F$ and all term orders \leq refining \mathbf{w} , $\text{LT}_{\leq}(f) = \text{in}_{\mathbf{w}}(f)$. Similarly, if \mathbf{w} defines a term order for an ideal I , then for all term orders \leq refining \mathbf{w} , $\text{LT}_{\leq}(I) = \text{in}_{\mathbf{w}}(I)$.

As seen in Lemma 3.3, for any finite set of polynomials F or any ideal I , and for any monomial order, there exists an equivalent monomial order defined by a system of weights.

Definition 5.6. Let $I \subseteq K[X]$ be an ideal. Let \mathbf{w} be a system of weights. The analytic Gröbner cone $C_{\mathbf{w}}(I)$ associated to \mathbf{w} and I is the set of all systems of weights \mathbf{w}' such that $\text{in}_{\mathbf{w}}(I) = \text{in}_{\mathbf{w}'}(I)$. The analytic Gröbner fan of I is the fan given by all the analytic Gröbner cones of I .

Proposition 5.7. If $\mathbf{w} \in \text{trop}(I)$, then $C_{\mathbf{w}}(I) \subset \text{trop}(I)$. In particular, the tropical variety associated to I is a subfan of the Gröbner fan of I .

PROOF. Whether a system of weights lies in the tropical variety only depends on the initial forms, and therefore applies identically to the cone. \square

Similar to the classical case, it allows to compute the tropical variety associated to I by traversing the Gröbner fan. The following properties are transpositions of corresponding facts in the classical setting, and describe the Gröbner fan.

Proposition 5.8. Let $I \subseteq K[X]$ be an ideal. Let \mathbf{w} be a system of weights, \mathbf{r} the convergence radii associated to \mathbf{w} , and \leq a term ordering refining \mathbf{w} . Let G be a reduced \mathbf{r} -local Gröbner basis of I (with \leq as tie-break). Then:

- (1) $\text{in}_{\mathbf{w}}(I) = \langle \text{in}_{\mathbf{w}}(g) : g \in G \rangle$;
- (2) for any system of weights \mathbf{w}_1 , $\mathbf{w}_1 \in C_{\mathbf{w}}(I)$ iff for all $g \in G$, $\text{in}_{\mathbf{w}}(g) = \text{in}_{\mathbf{w}_1}(g)$;
- (3) $C_{\mathbf{w}}(I)$ is the relative interior of a polyhedral convex cone;
- (4) the closure of $C_{\mathbf{w}}(I)$ in $\mathbb{R}_{<0} \times \mathbb{R}^n$ is the union of all cones $C_{\mathbf{w}'}(I)$ with $\forall f \in I$, $\text{in}_{\mathbf{w}}(f) = \text{in}_{\mathbf{w}'}(\text{in}_{\mathbf{w}}(f))$ (or equivalently $\forall g \in G$, $\text{in}_{\mathbf{w}}(g) = \text{in}_{\mathbf{w}'}(\text{in}_{\mathbf{w}}(g))$);
- (5) if \mathbf{w}_1 and \mathbf{w}_2 are two systems of weights such that $\overline{C_{\mathbf{w}_1}(I)} \cap \overline{C_{\mathbf{w}_2}(I)} \not\subseteq \{0\} \times \mathbb{R}^n$ and $C_{\mathbf{w}_1} \neq C_{\mathbf{w}_2}$, then there exists \mathbf{w}_3 such that $\overline{C_{\mathbf{w}_1}(I)} \cap \overline{C_{\mathbf{w}_2}(I)} = \overline{C_{\mathbf{w}_3}(I)}$, and it is a facet of both cones;
- (6) $C_{\mathbf{w}}(I)$ has maximal dimension $n+1$ if and only if \mathbf{w} defines a monomial order.

PROOF. For (1), let $f \in I$. Let $f_{\mathbf{r}}$ be the canonical image of f in $K\{X; \mathbf{r}\}$, it lies in $I_{\mathbf{r}}$. Since G is an \mathbf{r} -local GB of I , its embedding $G_{\mathbf{r}}$ in $I_{\mathbf{r}}$ is a Gröbner basis of $I_{\mathbf{r}}$ w.r.t. \leq . In particular, $f_{\mathbf{r}}$ reduces to 0 modulo $G_{\mathbf{r}}$, which in particular implies that there exists a finite sequence of reductions

$$f_{\mathbf{r}} \rightarrow f_{\mathbf{r},1} = f_{\mathbf{r}} - t_1 g_{\mathbf{r},j_1} \rightarrow \dots \rightarrow f_{\mathbf{r},k} = f_{\mathbf{r}} - \sum_{i=1}^k t_i g_{\mathbf{r},j_i}$$

with $g_{\mathbf{r},j} \in G_{\mathbf{r}}$ for all j , such that $\text{val}_{\mathbf{r}}(f_{\mathbf{r},k}) > \text{val}_{\mathbf{r}}(f_{\mathbf{r}})$ and $\text{val}_{\mathbf{r}}(f_{\mathbf{r},j}) = \text{val}_{\mathbf{r}}(f_{\mathbf{r}})$ for $j < k$. In particular, $\text{in}_{\mathbf{w}}(f_{\mathbf{r}}) = \sum_{i=1}^k t_i \text{in}_{\mathbf{w}}(g_{\mathbf{r},j_i})$. This is a polynomial equality, which translates into $\text{in}_{\mathbf{w}}(f) = \sum_{i=1}^k t_i \text{in}_{\mathbf{w}}(g_{j_i})$.

For (2), let \mathbf{w}_1 be a system of weight. The \Leftarrow direction follows from (1). Assume that $\mathbf{w}_1 \in C_{\mathbf{w}}(I)$, and consider the term order \leq_1 obtained by first considering the \mathbf{w}_1 -degree, then breaking ties using \leq . This term order refines \mathbf{w}_1 . Furthermore, $\text{LT}_{\leq_1}(I) = \text{LT}_{\leq_1}(\text{in}_{\mathbf{w}_1}(I)) = \text{LT}_{\leq}(\text{in}_{\mathbf{w}_1}(I))$, the last equality coming from the fact that the initial forms are \mathbf{w}_1 -homogeneous. Since $\text{in}_{\mathbf{w}_1}(I) = \text{in}_{\mathbf{w}}(I)$, we finally get that $\text{LT}_{\leq_1}(I) = \text{LT}_{\leq}(I)$. By Lemma 4.3, G is a reduced Gröbner basis w.r.t. \leq_1 .

From there, the proof is similar to that of [22, Prop. 3.1.17]. Let $g \in G$. Because G is reduced, $\text{LT}_{\leq_1}(g) = \text{LT}_{\leq}(g)$ and in particular this term lies in both the support of $\text{in}_{\mathbf{w}}(g)$ and $\text{in}_{\mathbf{w}_1}(g)$. Consider $\text{in}_{\mathbf{w}}(g) - \text{in}_{\mathbf{w}_1}(g)$, it lies in $\text{in}_{\mathbf{w}_1}(I)$ and its leading term w.r.t. \leq_1

cannot be $\text{LT}_{\leq_1}(g)$. Again since G is reduced, this implies that $\text{in}_{\mathbf{w}}(g) - \text{in}_{\mathbf{w}_1}(g) = 0$.

The next three properties are proved similarly to the classical case, by using the initial part conditions to cut half-spaces and define cones: see [22, Prop. 3.1.19] for (3), [22, Cor. 3.1.20] for (4) and [22, Prop. 3.1.21] for (5).

Finally, for (6), if \mathbf{w} defines a monomial order for I , by considering the finite set $\text{Supp}(G)$, there exists a small enough $\epsilon \in \mathbb{R}_{>0}$ such that for all $\mathbf{u} \in \mathbb{R}_{<0} \times \mathbb{R}^n$, and for all $t_1, t_2 \in \text{Supp}(G)$, $\deg_{\mathbf{w}_1}(t_1) < \deg_{\mathbf{w}_2}(t_2) \iff \deg_{\mathbf{w}_1+\epsilon\mathbf{u}}(t_1) < \deg_{\mathbf{w}_2+\epsilon\mathbf{u}}(t_2)$. Therefore $C_{\mathbf{w}}(I)$ contains a ball of radius ϵ , so it is open and it must have maximal dimension. Conversely, if \mathbf{w} does not define a monomial order, there exists a $g \in G$ such that $\text{in}_{\mathbf{w}}(g)$ has at least two terms t_1 and t_2 , so \mathbf{w} lies on the hyperplane $H = \{\text{val}_{\mathbf{w}}(t_1) = \text{val}_{\mathbf{w}}(t_2)\}$. For all $\mathbf{w}' \in C_{\mathbf{w}}(I)$, $\text{in}_{\mathbf{w}'}(g) = \text{in}_{\mathbf{w}}(g)$, so $C_{\mathbf{w}}(I) \subseteq H$, and it cannot have maximal dimension. \square

Since we know that I has finitely many sets of leading terms by Th. 4.2, the analytical Gröbner fan of I has only finitely many cones of maximal dimension, and therefore it is finite. If furthermore I is homogeneous, then I admits a universal Gröbner basis G containing a reduced Gröbner basis for all orders, computable using Algorithm 2. In that case, by Prop. 3.8, the vertices of $\mathcal{N}(G)$ are in one-to-one correspondence with the equivalence classes of term orders with respect to G , and by Lemma 3.2, this allows to compute all the equivalence classes of term orders with respect to I . From there one can compute the cones of maximal dimension in the analytic Gröbner fan of I , and finally all cones in the fan.

Then, for each cone C in the fan, one can pick a system of weights \mathbf{w} in C . From the property (1), $\text{in}_{\mathbf{w}}(I)$ is generated by the initial parts of elements of G . Finally, we can decide whether $\text{in}_{\mathbf{w}}(I)$ contains a monomial by computing a basis of $(\text{in}_{\mathbf{w}}(I) : (x_1 \cdots x_n)^\infty)$, using the algorithms in [7].

Just like in the classical case, both with and without valuation, this algorithm is not the most efficient, because the Gröbner fan can be significantly larger than the tropical variety. In [22] and [17], better algorithms have been presented for ideals in $K[[T]][X]$, and generalized to p -adic fields by lifting back to that case. Those algorithms still traverse the Gröbner fan, but not in an exhaustive way, and rely on Buchberger's algorithm with Mora reductions for computing bases in the cones.

The conclusion of this section is that the key properties of the Gröbner fan in $K[[T]][X]$ are shared across polynomial rings over a valued ring or field, by way of Tate completions, and Buchberger's algorithm with weak normal form allows to compute generators in a similar way. Thus, it offers an alternative point of view on existing algorithms using liftings to reduce the problem to $K[[T]][X]$, instead performing the computations directly in the desired ring or field.

We expect that the other optimized algorithms for computing tropical varieties in $K[[T]][X]$ similarly transpose to the general setting.

Remark 5.9. The requirement that the ideal is homogeneous can be relaxed into requiring that the ideal admits a reduced Gröbner basis for all orders. This requirement is also present in the aforementioned works in $K[[T]][X]$.




Figure 1: An example of a polyhedron with \mathcal{P} the convex hull of the $s_i + \mathbb{R}_{<0}^2$'s in \mathbb{R}^2 .

6 TATE ALGEBRAS ON POLYHEDRAL SUBDOMAINS

As of now, we have studied Tate algebras using convergence conditions of the following types: (1) on a polydisk defined by log-radii $\mathbf{s} \in \mathbb{Q}^n$; (2) convergence everywhere, i.e. the algebra $K[X]$; (3) on all polydisks $\mathbf{r} \leq \mathbf{s}$ (overconvergence).

To build up the tools for rigid geometry or tropical analytic geometry as in [21], we need more general convergence conditions such as convergence on an annulus (e.g. converging for all $\mathbf{X} \in \mathbb{Q}_p^n$ such that $\forall i, a_i \leq \text{val}(x_i) \leq b_i$ for some $a_i, b_i \in \mathbb{R}$) or converging on a polyhedron.

Following [21, Definition 6.3] but restricting to the case of power series instead of Laurent series, we define the following ring of functions $K\{X; P\}$ for P the rational points of a polyhedron. This corresponds to a special case of the affinoid algebra defined by a polyhedral subdomain in [21].

Definition 6.1. Let $\mathcal{P} \subset \mathbb{R}^n$ be a polyhedron with vertices in \mathbb{Q}^n and admitting only $\sigma = \mathbb{R}_{<0}^n$ as cone of unbounded direction. Let $P = \mathcal{P} \cap \mathbb{Q}^n$. We define the ring

$$K\{X; P\} := \left\{ \sum_{\alpha \in \mathbb{N}^n} a_\alpha X^\alpha : \begin{array}{l} a_\alpha \in K \\ \forall \mathbf{r} \in P, \text{val}_{\mathbf{r}}(a_\alpha X^\alpha) \rightarrow +\infty \end{array} \right\}.$$

The series in $K\{X; P\}$ are exactly the series converging on all polydisks with radius given by the weights which are points of P . We give an example in Figure 1.

Proposition 6.2. If \mathcal{P} is the convex hull of the $s_1 + \mathbb{R}_{<0}^n, \dots, s_l + \mathbb{R}_{<0}^n$ then

$$K\{X; P\} = \bigcap_{i=1}^l K\{X; s_i\}.$$

PROOF. The \subset inclusion is clear. For the converse inclusion, we remark that if $\mathbf{r} \in P$, there are some $\lambda_i \geq 0$ such that $\sum_{i=1}^l \lambda_i = 1$ and $\sum_{i=1}^l \lambda_i s_i \geq \mathbf{r}$. Consequently, for any term $c_\alpha X^\alpha \in K[X]$,

$$\sum_{i=1}^l \lambda_i \text{val}_{s_i}(c_\alpha X^\alpha) \leq \text{val}_{\mathbf{r}}(c_\alpha X^\alpha).$$

If $f = \sum_{\alpha \in \mathbb{N}^n} c_\alpha X^\alpha \in \bigcap_{i=1}^l K\{X; s_i\}$ then for all i , $\text{val}_{s_i}(c_\alpha X^\alpha) \rightarrow +\infty$. Thus, $\text{val}_{\mathbf{r}}(c_\alpha X^\alpha) \rightarrow +\infty$. Hence, we get that, if $f \in \bigcap_{i=1}^l K\{X; s_i\}$, then $f \in K\{X; P\}$ which concludes the proof. \square

6.1 Local Gröbner bases

In this section, we explain how if \mathcal{P} is such a polyhedron, $P = \mathcal{P} \cap \mathbb{Q}^n$, I an ideal in $K\{X; P\}$, and $\mathbf{r} \in P$, it is possible to compute an \mathbf{r} -local Gröbner basis of I comprised only of elements of $K\{X; P\}$. First, we adapt the notion of écart from [7, §6.1].

Definition 6.3. Let $f \in K\{X; P\}$, and $\mathbf{r}, \mathbf{s} \in P$. We define the (\mathbf{s}, \mathbf{r}) -degree of f as $\deg_{\mathbf{s}, \mathbf{r}}(f) = \max_{\alpha \in \text{Supp}_s(f)} (\mathbf{s} - \mathbf{r}) \cdot \alpha$, and the écart of first and second type of f as

$$\begin{aligned}\hat{\text{Ecart}}_{\mathbf{s}, \mathbf{r}, 0}(f) &:= \text{val}_{\mathbf{s}}(\text{LT}_{\mathbf{r}}(f)) - \text{val}_{\mathbf{s}}(f), \\ \hat{\text{Ecart}}_{\mathbf{s}, \mathbf{r}, 1}(f) &:= \deg_{\mathbf{s}, \mathbf{r}}(f) - \deg_{\mathbf{s}, \mathbf{r}}(\text{LT}_{\mathbf{r}}(f))\end{aligned}$$

Lemma 6.4 (6.3 in [7]). $\forall f \in K\{X; P\}, \forall i = 0, 1, \hat{\text{Ecart}}_{\mathbf{s}, \mathbf{r}, i}(f) \geq 0$.

We adapt Mora's overconvergent Weak Normal Form algorithm of [18] to computations in $K\{X; \mathbf{r}\}$ for series in $K\{X; P\}$, as in [7].

Algorithm 3 WNF(f, g, P, \mathbf{r}), Mora's WNF for local computations in polyhedral subdomains

Input: $f, g_1, \dots, g_s \in K\{X; P\}$ where \mathcal{P} is the convex hull of the $s_1 + \mathbb{R}_{\leq 0}^n, \dots, s_t + \mathbb{R}_{\leq 0}^n$ in \mathbb{R}^n for some s_i 's, $P = \mathcal{P} \cap \mathbb{Q}^n$ and $\mathbf{r} \in P$.

Output: $h \in K\{X; P\}$ such that:

- for some $\mu, u_1, \dots, u_s \in K\{X; P\}, \mu f = \sum u_i g_i + h$
- either $h = 0$ or $\text{LT}_{\mathbf{r}}(h)$ is divisible by no $\text{LT}_{\mathbf{r}}(g_i)$
- μ is invertible in $K\{X; \mathbf{r}\}$
- $\text{LT}_{\mathbf{r}}(u_i g_i) \leq \text{LT}_{\mathbf{r}}(f)$

```

1:  $h := f$ ;
2:  $T := (g_1, \dots, g_s)$ ;
3: while  $h \neq 0$  and  $T_h := \{g \in T, \text{LT}_{\mathbf{r}}(g) \mid \text{LT}_{\mathbf{r}}(h)\} \neq \emptyset$  do
4:   take  $g \in T_h$  minimizing  $\hat{\text{Ecart}}_{s_1, \mathbf{r}, 0}(g)$  first, then  $\hat{\text{Ecart}}_{s_1, \mathbf{r}, 1}(g)$ ,
   then  $\hat{\text{Ecart}}_{s_2, \mathbf{r}, 0}(g), \dots$ , and finally  $\hat{\text{Ecart}}_{s_t, \mathbf{r}, 1}(g)$ ;
5:   if for any  $j, k, \hat{\text{Ecart}}_{s_j, \mathbf{r}, k}(g) > \hat{\text{Ecart}}_{s_j, \mathbf{r}, k}(h)$ , then
6:      $T := T \cup \{h\}$ ;
7:    $h := \text{S-Poly}(h, g)$ ;
8: return  $h$ ;
```

Correctness comes from the following.

Lemma 6.5 (6.4 in [7]). *If $g \in T_{h_m}$ is such that:*

- $\hat{\text{Ecart}}_{s_i, \mathbf{r}, 0}(g) \leq \hat{\text{Ecart}}_{s_i, \mathbf{r}, 0}(h_m)$,
- $\hat{\text{Ecart}}_{s_i, \mathbf{r}, 1}(g) \leq \hat{\text{Ecart}}_{s_i, \mathbf{r}, 1}(h_m)$,

and if $t = \text{LT}_{\mathbf{r}}(h_m)/\text{LT}_{\mathbf{r}}(g)$ and $h_{m+1} = h_m - tg$, then

$$\text{val}_{s_i}(h_{m+1}) \geq \text{val}_{s_i}(h_m).$$

Moreover, in case of equality, then

$$\deg_{s_i, \mathbf{r}}(h_{m+1}) \leq \deg_{s_i, \mathbf{r}}(h_m).$$

Proposition 6.6. *Algorithm 3 either terminates in a finite number of steps, or $\text{LT}_{\mathbf{r}}(h)$ and all the $\text{LT}_{s_i}(h)$'s converge to 0.*

PROOF. The proof follows exactly the same lines as that of [7, Prop. 6.5]. The only difference is the definition of the extended leading terms: for $h \in K\{X; P\}$ we define

$$\text{LTE}(h) := \prod_{i=1}^t U_i^{d_{i,1}} \hat{\text{Ecart}}_{s_i, \mathbf{r}, 0}(h) \prod_{i=1}^t V_i^{d_{i,2}} \hat{\text{Ecart}}_{s_i, \mathbf{r}, 1}(h) \text{LT}(h)$$

in $K[X, U_1, \dots, U_t, V_1, \dots, V_t]$, for some suitable $d_{i,j} \in \mathbb{N}$ making everything nonnegative integers. \square

Using Algorithm 3 as a replacement for the reduction procedure in Buchberger's algorithm of [7, Sec. 5], one can compute an \mathbf{r} -local GB of an ideal of $K\{X; P\}$.

6.2 Conjectures

The present work is only scratching the surface of the study of ideals in $K\{X; P\}$. As an example, we conclude with some questions regarding statements which hold in $K[X]$ or $K\{X; \mathbf{r}\}$, but are unclear in $K\{X; P\}$.

Conjecture 6.7. *If $I \subset K\{X; P\}$ is an ideal, then the set $\text{Terms}_P(I) = \{\text{LT}_{\mathbf{r}}(I), \text{ for } \mathbf{r} \in P\}$ is finite.*

We may remark that this conjecture is true for principal ideals.

Proposition 6.8. *If $f \in K\{X; P\}$, $\text{Terms}_P(\langle f \rangle)$ is finite.*

PROOF. Let $s_1, \dots, s_l \in \mathbb{Q}^n$ be the extremal points of \mathcal{P} . Then \mathcal{P} is the convex hull of the $s_i + \mathbb{R}_{\leq 0}^n$ (and $P = \mathcal{P} \cap \mathbb{Q}^n$). Let $S(f)$ be the set of all $(\text{val}_{s_1}(c_{\alpha} X^{\alpha}), \dots, \text{val}_{s_l}(c_{\alpha} X^{\alpha}), \alpha_1, \dots, \alpha_n)$ for $c_{\alpha} X^{\alpha} \in \text{Supp}(f)$. Up to scalar multiplication of f , we can assume that $S(f) \subset \left(\frac{1}{D} \mathbb{N}^l\right) \times \mathbb{N}^n$ for some $D \in \mathbb{N}$. Hence, $S(f)$ only has a finite number of minimal elements for the product order on $\left(\frac{1}{D} \mathbb{N}^l\right) \times \mathbb{N}^n$. Let us denote them by Σ .

Let $\mathbf{r} \in P$. We prove that if $t = \text{LT}_{\mathbf{r}}(f)$, then $t \in \Sigma$. Let us write $t = c_{\alpha} X^{\alpha}$, and let $u = c_{\beta} X^{\beta}$ be a term of f , belonging to Σ and smaller than t for the product order: $\text{val}_{s_i}(u) \leq \text{val}_{s_i}(t)$ for all $i \leq l$, and $\beta_i \leq \alpha_i$ for all $i \leq n$.

Since $\mathbf{r} \in P \subset \mathcal{P}$ and \mathcal{P} is the convex hull of the $s_i + \mathbb{R}_{\leq 0}^n$, there are some $\lambda_i \geq 0$ such that $\sum_{i=1}^l \lambda_i = 1$ and $\sum_{i=1}^l \lambda_i s_i \geq \mathbf{r}$. Then:

$$\begin{aligned}\text{val}_{\mathbf{r}}(u) &= \text{val}(c_{\beta}) - \mathbf{r} \cdot \beta, \\ &= \sum_{i=1}^l \lambda_i (\text{val}(c_{\beta}) - s_i \cdot \beta) + \left(\sum_{i=1}^l \lambda_i s_i - \mathbf{r} \right) \cdot \beta, \\ &= \sum_{i=1}^l \lambda_i \text{val}_{s_i}(u) + \left(\sum_{i=1}^l \lambda_i s_i - \mathbf{r} \right) \cdot \beta.\end{aligned}$$

Since for all $i \leq l$, $\text{val}_{s_i}(u) \leq \text{val}_{s_i}(t)$ and for all $i \leq n$, $\beta_i \leq \alpha_i$ and $\sum_{i=1}^l \lambda_i s_i \geq \mathbf{r}$, then

$$\text{val}_{\mathbf{r}}(u) \leq \sum_{i=1}^l \lambda_i \text{val}_{s_i}(t) + \left(\sum_{i=1}^l \lambda_i s_i - \mathbf{r} \right) \cdot \alpha \leq \text{val}_{\mathbf{r}}(t).$$

Consequently, $\text{val}_{\mathbf{r}}(u) \leq \text{val}_{\mathbf{r}}(t)$ and $\text{val}_{\mathbf{r}}(t)$ is the minimum of the $\text{val}_{\mathbf{r}}(v)$ for v a term of f . Hence $\text{val}_{\mathbf{r}}(u) = \text{val}_{\mathbf{r}}(t)$, and thus all the above inequalities are equalities. It then means that $u = t$ and $\text{LT}_{\mathbf{r}}(f) \in \Sigma$, as claimed. Since Σ is finite, and $\text{Terms}_P(\langle f \rangle) = \{\text{LT}_{\mathbf{r}}(f), \mathbf{r} \in P\}$ the proposition is proved. \square

Our end goal is the following stronger conjecture.

Conjecture 6.9. *If $I \subset K\{X; P\}$ is an ideal, then there is a UAGB of I i.e. a finite set $G \subset I$ such that G is an \mathbf{r} -local GB of I for any $\mathbf{r} \in P$.*

We expect zero-dimensional ideals of $K\{X; P\}$ to be polynomial ideals and the conjecture is clearly true for all polynomial ideals.

Remark 6.10. The main issue encountered in attempting to prove Conjectures 6.7 and 6.9 is the fact that for some $\mathbf{r} \in P$, there exists a reduced Gröbner basis of $I_{\mathbf{r}}$, but it usually does not belong to I and thus is not converging for other points of P . Hence we can not rely on Lemma 4.3. This issue was overcome for polynomials using homogenization (Section 4.2 and [7, Sec. 7]), but this technique does not directly translate to the case of series.

REFERENCES

- [1] Allamigeon X., Benchimol P. N., Gaubert S., Joswig M. Log-Barrier Interior Point Methods Are Not Strongly Polynomial SIAM Journal on Applied Algebra and Geometry, 2018, vol 2
- [2] Bogart T., Jensen A. N., Speyer D., Sturmfels B., Thomas R. R. Computing Tropical Varieties J. Symb. Comput. 42 (2007), no. 1-2
- [3] Buchberger B., Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem nulldimensionalen Polynomideal (An Algorithm for Finding the Basis Elements in the Residue Class Ring Modulo a Zero Dimensional Polynomial Ideal), English translation in J. of Symbolic Computation, Special Issue on Logic, Mathematics, and Computer Science: Interactions. Vol. 41, Number 3-4, Pages 475–511, 2006
- [4] Caruso X., Vaccon T., Verron T., Gröbner bases over Tate algebras, in Proceedings: ISSAC 2019, Beijing, China.
- [5] Caruso X., Vaccon T., Verron T., Signature-based algorithms for Gröbner bases over Tate algebras, in Proceedings: ISSAC 2020, Kalamata, Greece.
- [6] Caruso X., Vaccon T., Verron T., On FGLM Algorithms With Tate Algebras, in Proceedings: ISSAC 2021, Saint-Petersburg, Russia.
- [7] Caruso X., Vaccon T., Verron T., On Polynomial Ideals and Overconvergence in Tate Algebras, in Proceedings: ISSAC 2022, Lille, France.
- [8] Chan A., MacLagan D., Gröbner bases over fields with valuations, Math. Comp. 88 (2019), 467-483.
- [9] Collart S., Kalkbrenner M., Mall D., Converting bases with the Gröbner walk., J. Symbolic Comp. 6 (1997), 209–217.
- [10] Cox D., Little J., O’Shea D., *Ideals, varieties, and algorithms*. Undergraduate Texts in Mathematics. Springer, Cham, fourth edition, 2015.
- [11] Cox D., Little J., O’Shea D., *Using Algebraic Geometry*. Graduate Texts in Mathematics, Volume 185, Springer Science & Business Media, 2005.
- [12] Gritzmann P., Sturmfels B. *Minkowski Addition of Polytopes: Computational Complexity and Application to Gröbner Bases*. SIAM J. Disc. Math., 1993.
- [13] Fukuda K., Jensen A. N., Thomas R. R., Computing Gröbner fans, Math. Comp. 76 (2007)
- [14] Greuel G.-M., Pfister G., A Singular Introduction to Commutative Algebra, Springer-Verlag Berlin Heidelberg 2008.
- [15] Jensen, A.N., Gfan, a software system for Gröbner fans and tropical varieties, Available at <http://home.imf.au.dk/jensen/software/gfan/gfan.html>
- [16] Mikhalkin G., Enumerative Tropical Algebraic Geometry In \mathbb{R}^2 , Journal of the American Aathematical Society, Volume 18, Number 2, 2005
- [17] Markwig T., Ren Y., Computing Tropical Varieties Over Fields with Valuation, Found Comput Math 20, 783–800 (2020).
- [18] Mora T., An Algorithm to Compute the Equations of Tangent Cones, in: Proceedings EUROCAM 82, Lecture Notes in Comput. Sci. (1982).
- [19] Mora T., Robbiano L., The Gröbner fan of an ideal, J. Symbolic Comput. 6 (1988), no. 2-3.
- [20] Ostrowski A.M., On Multiplication and Factorization of Polynomials, I. Lexicographical Orderings and Extreme Aggregates of Terms, Aequationes Mathematicae, Volume 13, 1975.
- [21] Rabinoff J., Tropical analytic geometry, Newton polygons, and tropical intersections, Advances in Mathematics 229(6), 2010.
- [22] Ren Y., Tropical geometry in Singular, Doctoral dissertation, Technische Universität Kaiserslautern, 2015.
- [23] SageMath, the Sage Mathematics Software System (Version 9.2), The Sage Development Team, 2020, <http://www.sagemath.org>
- [24] Sturmfels B., Gröbner Bases and Convex Polytopes, American Mathematical Society, Univ. Lectures Series, No 8, Providence, Rhode Island, 1996.
- [25] Tate J., Rigid analytic spaces, Inventiones Mathematicae 12, 1971, 257–289
- [26] Vaccon T., Matrix-F5 Algorithms and Tropical Gröbner Bases Computation, in Proceedings: ISSAC 2015, Bath, UK. Extended version in the J. of Symbolic Computation, 2017.
- [27] Vaccon T., Yokoyama K., A Tropical F5 algorithm, in Proceedings: ISSAC 2017, Kaiserslautern, Germany.
- [28] Vaccon T., Verron T., Yokoyama K., On Affine Tropical F5 algorithm, in Proceedings: ISSAC 2018, New York, USA. Extended version in the J. of Symbolic Computation, 2021.